



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Hot Electron Diagnostics using X-rays and Cerenkov Radiation

J. Stein, E. E. Fill, G. Pretzler, F. Brandl, J. Kuba,
D. Habs

January 12, 2004

Ultrashort High-Energy Radiation and Matter
Varenna, Italy
October 7, 2003 through October 10, 2003

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

Ultrashort High -Energy Radiation and Matter, Varenna, Oct. 7 -10, 2003

Hot electron diagnostics using X -rays and Cerenkov radiation

J. Stein ^{1,2}, E. Fill ¹, G. Pretzler ³, F. Brandl ³, J. Kuba ⁴ and D. Habs ²

¹*Max-Planck-Institut für Quantenoptik, D -85748 Garching, Germany*

²*Ludwig-Maximilians-Universität München, D -85748 Garching, Germany*

³*Universität Düsseldorf, D -40225 Düsseldorf, Germany*

⁴*Lawrence Livermore National Laboratory, Livermore CA 94550, USA*

Abstract: The propagation of laser -generated hot electron s through matter and across narrow vacuum gaps is studied. We use the ATLAS titanium -sapphire laser of Max -Planck-Institut für Quantenoptik to irradiate 10 μm to 100 μm thick copper foils at intensities up to 10¹⁹ W/cm², generating electrons with temperatures in the MeV -range. After propagating through the target the electrons are detected via Cerenkov radiation generated in a suitable medium and by hard X -ray emitted from an X -ray "fluor". In some experiments a plastic scintillator was used to monitor the electrons. These diagnostics allow to characterize the electrons with respect to their energy, number and directionality. We also investigate the propagation of the hot electrons across narrow vacuum gaps, with a width ranging from several 100 μm down to 25 μm. The effect of self -generated fields ¹ in preventing electrons to cross the gap is demonstrated. Implications of these experiments with respect to pumping of X -ray lasers ², isochoric heating by X -rays³ and developing optics for 4th -generation light sources will be discussed.

¹E.E.Fill, Phys. Plasmas 8(2001)4613

²G.Pretzler, Th. Schlegel, E.Fill, and D.Eder, Phys.Rev.E62,(2000)5618.

³G.Dyer, R.Shepherd, J.Kuba, E.Fill, A.Wootton, P.Patel, D.Price and T.Ditmire, J.Mod. Opt., in print.